

BacterLab Division



Issue date: 04/08/2025

Version: 01.2025

SURFACE SAMPLE COLLECTION MEDIUM

BacterContact™ Tryptic Soy Agar + Penase + Cephalosporinase + Neutralizing

Ready-to-use medium on 60mm plates for environmental and personnel monitoring with inactivation of β -lactam antibiotics and disinfectants.

Code: 4109035



1. INTENDED USE

BacterContact™ Tryptic Soy Agar + Penase + Cephalosporinase + Neutralizing is a general purpose medium in RODAC (Replicate Organisms Detection and Counting) plates used for environmental and personnel monitoring with inactivation of β -lactam antibiotics (penicillins and cephalosporins) and disinfectants.

The packaging with semi-permeable Cellophane film helps balance the humidity of the environment during storage.

2. PRINCIPLES

Casein peptone and soy peptone provide amino acids, nitrogen, carbon, minerals, vitamins and other nutrients which support the growth of microorganism. Sodium chloride maintains the osmotic balance of the medium. Agar is the solidifying agent. Histidine inactivates aldehydes. Lecithin neutralizes quaternary ammonium compounds. Polysorbate 80 (Tween 80) is effective against phenolic compounds and mercurial derivatives. Sodium thiosulfate neutralizes halogen compounds. Penase is a preparation of Penicillinase and with Cephalosporinase inactivate β -lactam antibiotics (penicillins and cephalosporins).

- 1 Levy Unit (LU) is defined as the amount of penicillinase that inactivates 59.3 IU of Penicillin G per hour at pH 7.0 at 25°C.
- International Unit (IU) is defined as the amount of enzyme needed to hydrolyze 1 μ mole of Penicillin G (Penicillinase) or 1 μ mole of Cephalosporin C (Cephalosporinase) per minutes at pH 7.0 at 25°C.

3. TYPICAL COMPOSITION

For 1 liter of medium (refractive)

Casein Peptone	15,0 g
Soy Peptone	5,0 g
Sodium Chloride	5,0 g
Agar	15,0 g
Histidine	1,0 g
Lecithin	0,7 g
Polysorbate 80	5,0 g
Sodium Thiosulfate	0,5 g
Penase	1,0 mL
Cephalosporinase	1,0 mL

pH of the ready-to-use medium at 25°C: $7,3 \pm 0,2$

4. PREPARATION

The environmental plates are ready-to-use, no preparation required.

5. INSTRUCTIONS FOR USE

Preparation:

- Prepare a test diagram for the areas that are to be tested and label plates with the corresponding location identification. Ensure that the label cannot be readily wiped off or removed. RODAC plates are prepared so that the agar surface is convex for sampling flat surfaces. Prior to sampling, the plates should be warmed to room temperature in the plastic sleeve for approximately 15 – 20 minutes with agar up and the lid down. Remove the quantity of plates from the sleeve that are required for testing. The location/ site identification should be written on the base (agar portion, not the lid) of the plate.
- RODAC plates are recommended for use in air sampling equipment as well as for surface sampling. Selected surfaces are sampled by firmly pressing the agar medium against the test area. Plates are intended for personnel hygiene monitoring (clothing, gloves or hands) as well.

Sampling:

- While wearing gloves, remove the lid from the plate with one hand. With the other hand, hold the base (agar portion of the plate) with thumb and middle finger. Use the index finger to gently press the plate on the test surface. Make sure the entire agar surface touches the test surface. Do not move the plate laterally while sampling as it will spread contaminants making enumeration difficult. Place the lid back on the plate and tape closed. Be sure to clean the test area after sampling to remove any residual growth media remaining on the surface.
- For detection of bacteria incubate the plates at 30 – 35°C for 18 – 72 hours.
- For detection of yeasts and moulds incubate at 20 – 25°C for 2 – 7 days.

6. RESULTS

Observe for the formation of fungal colonies exhibiting typical microscopic and colonial morphology. Record the number of CFU per plate. Colonies should be further isolated and identified with appropriate procedures.

7. QUALITY CONTROL

BacterLab ensures the quality of each product batch by testing with ATCC reference strains.

Reference strains	Incubation conditions	Expected results
<i>Candida albicans</i> ATCC 10231	Incubate for 48 – 72 hours at 20 – 25°C	Good growth
<i>Aspergillus brasiliensis</i> ATCC 16404		
<i>Staphylococcus aureus</i> ATCC 25923	Incubate for 18 – 24 hours at 30 – 35°C	
<i>Pseudomonas aeruginosa</i> ATCC 9027		
<i>Escherichia coli</i> ATCC 25922		

8. STORAGE AND TRANSPORT CONDITIONS

- Storage: 2 – 8°C.
- Transportation: Ambient temperature.

9. PACKAGING

- Packaging: 10 plates/ box or as per customer request.

10. SHELF LIFE

- Expiration Date: 06 months from the manufacturing date.

11. BIBLIOGRAPHY

- Swanson, K.J., F.F. Busta, E.H. Peterson, and M.G. Johnson (1992). Colony Count Methods, p. 75-95.
- USP 33 – NF 28 (2011) Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. The United States Pharmacopeial Convention, Rockville, MD. USA.
- USP 33 – NF 28 (2011) Microbiological evaluation of clean rooms. The United States Pharmacopeial Convention, Rockville, MD. USA.
- European Pharmacopeia 7.0 (2011) 7th ed. Chapters 2.6.13. Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. Council of Europe Strasbourg, France.
- Solabia Group. *Tryptic Soy Agar*. Biokar Diagnostics.